arXiv:1010.0089v1 [hep-ph] 1 Oct 2010

Nikhef-2010-033

Extracting γ and Penguin Topologies through CP Violation in $B_s^0 \to J/\psi K_{\rm S}$

Kristof De Bruyn, Robert Fleischer and Patrick Koppenburg

Nikhef, Science Park 105, NL-1098 XG Amsterdam, The Netherlands

Abstract

The $B_s^0 \to J/\psi K_{\rm S}$ decay has recently been observed by the CDF collaboration and will be of interest for the LHCb experiment. This channel will offer a new tool to extract the angle γ of the unitarity triangle and to control doubly Cabibbo-suppressed penguin corrections to the determination of $\sin 2\beta$ from the well-known $B_d^0 \to J/\psi K_{\rm S}$ mode with the help of the U-spin symmetry of strong interactions. While any competitive determination of γ is interesting, the latter aspect is particularly relevant as LHCb will enter a territory of precision which makes the control of doubly Cabibbo-suppressed Standard-Model corrections mandatory. Using the data from CDF and the e^+e^- B factories as a guideline, we explore the sensitivity for γ and the penguin parameters and point out that the B_s^0–$\bar{B}_s^0$ mixing phase ϕ_s, which is only about -2° in the Standard Model but may be enhanced through new physics, is a key parameter for these analyses. We find that the mixing-induced CP violation $S(B_s^0 \to J/\psi K_{\rm S})$ shows an interesting correlation with $\sin\phi_s$, which serves as a target region for the first measurement of this observable at LHCb.

September 2010

1 Introduction

As the LHCb experiment at CERN's Large Hadron Collider (LHC) has just started to take first physics data, we are entering a new territory in precision flavour physics. A particularly exciting feature of the LHCb research programme is the exploration of decays of B_s mesons. Among the various B_s decays with a promising physics potential is the $B_s^0 \to J/\psi K_{\mathrm{S}}$ mode, which has recently been observed by the CDF collaboration at the Tevatron [1].

In Ref. [2], it was pointed out that the CP-violating asymmetries of this channel offer a probe for extracting the angle γ of the unitarity triangle (UT). Moreover, it was noted that also penguin topologies can be determined, which limit the theoretical precision of the measurement of $\sin 2\beta$, where β is another UT angle, by means of the mixing-induced CP violation in the "golden" decay $B_d^0 \to J/\psi K_{\mathrm{S}}$. In this strategy, the U-spin flavour symmetry of strong interactions is used, which is a subgroup of $SU(3)_{\mathrm{F}}$ relating down and strange quarks to each other, and allows us to relate the hadronic parameters of the $B_s^0 \to J/\psi K_{\mathrm{S}}$ and $B_d^0 \to J/\psi K_{\mathrm{S}}$ modes to one another. In Fig. 1, we show the decay topologies of the $B_s^0 \to J/\psi K_{\mathrm{S}}$ channel; interchanging all down and strange quarks, we obtain the $B_d^0 \to J/\psi K_{\mathrm{S}}$ topologies.



Figure 1: Decay topologies contributing to the $B_s^0 \to J/\psi K_{\mathrm{S}}$ channel: "tree" (left) and "penguin" (right) diagrams.

In view of the start of LHCb, we would like to have a closer look at the prospects for analysing the $B_s^0 \to J/\psi K_{\mathrm{S}}$ channel at this experiment. Here we will give a special emphasis to the extraction of the hadronic penguin parameters and the control of the corresponding uncertainty in the measurement of $\sin 2\beta$ from $B_d^0 \to J/\psi K_{\mathrm{S}}$. In Ref. [3], using data on the $B_d^0 \to J/\psi \pi^0$ channel (see also [4]), it was pointed out that such effects tend to lower a tension in the fit of the UT between measurements of its side $R_b \propto |V_{ub}/V_{cb}|$ and $\sin 2\beta$. Following these results, we will explore the prospects for such an analysis by means of the $B_s^0 \to J/\psi K_{\mathrm{S}}$ channel as an analysis of $B_d^0 \to J/\psi \pi^0$ is considered to be more challenging at LHCb.

Due to the low branching fraction of the $B_s^0 \to J/\psi K_{\mathrm{S}}$ mode, LHCb might be able to get only very crude first information on the corresponding CP asymmetries by the end of 2011, even though a signal should be clearly visible by then. In order to explore the full potential of this decay, we will therefore assume two milestones for our feasibility study:

- an integrated luminosity of 6 fb^{-1}, which the LHCb experiment could realistically collect by the end of the second data taking period of 2014–15 (with the LHC having reached the design centre-of-mass energy $\sqrt{s} = 14$ TeV);
- an integrated luminosity of 100 fb^{-1}, which is the current target for a subsequent LHCb upgrade.

For the measurement of $\sin 2\beta$ through $B_d^0 \to J/\psi K_{\mathrm{S}}$, $(\sin 2\beta)_{J/\psi K_{\mathrm{S}}}$, LHCb expects precisions of 0.022 with 2 fb^{-1} [5], and 0.003–0.010 for 100 fb^{-1}, depending on to be determined systematic errors [6]. In view of these impressive numbers, we have to take penguin effects into account in order to match the experimental with the theoretical precision. Such an analysis may eventually also allow us to pin down new-physics (NP) effects in B_d^0–$\bar{B}_d^0$ mixing; the current data still leave space for NP contributions as large as about 50% of the SM contribution, with a CP-violating NP phase below 10° (see, for instance, Ref. [7]).

The outline of this paper is as follows: in Section 2, we discuss the strategy to extract γ and the penguin parameters from the $B_{d,s} \to J/\psi K_{\mathrm{S}}$ modes. In Section 3, we have a look at the picture for the relevant observables emerging from the current data, and, after introducing the framework for the LHCb feasibility study in Section 4, discuss then in Section 5 the prospects for the resulting determination of γ. In Section 6, we focus on the extraction of the penguin parameters, and discuss then the control of their impact on the measurement of $(\sin 2\beta)_{J/\psi K_{\mathrm{S}}}$ in Section 7. Finally, we summarise our conclusions and give a brief outlook in Section 8.

2 The Basic Strategy

The $B_s^0 \to J/\psi K_{\mathrm{S}}$ transition originates from a $\bar{b} \to \bar{c}c\bar{d}$ quark-level process and has a decay amplitude of the following structure:

$$A(B_s^0 \to J/\psi\, K_{\mathrm{S}}) = \lambda_c^{(d)} \left[A_{\mathrm{T}}^{(c)} + A_{\mathrm{P}}^{(c)}\right] + \lambda_u^{(d)} A_{\mathrm{P}}^{(u)} + \lambda_t^{(d)} A_{\mathrm{P}}^{(t)}\,, \tag{1}$$

where $A_{\mathrm{T}}^{(c)}$ and $A_{\mathrm{P}}^{(j)}$ denote strong amplitudes that are related to the "tree" and "penguin" topologies (with internal $j \in \{u, c, t\}$ quarks), respectively, which are illustrated in Fig. 1. On the other hand, the

$$\lambda_q^{(d)} \equiv V_{qd} V_{qb}^* \tag{2}$$

factors contain the relevant combinations of elements of the Cabibbo–Kobayashi–Maskawa (CKM) matrix. If we make use of the unitarity of the CKM matrix and apply the Wolfenstein parametrisation [8] as generalised in Ref. [9], we can write (1) as

$$A(B_s^0 \to J/\psi\, K_{\mathrm{S}}) = -\lambda\, \mathcal{A}\left[1 - a e^{i\theta} e^{i\gamma}\right]. \tag{3}$$

Here

$$\mathcal{A} \equiv \lambda^2 A \left[A_{\mathrm{T}}^{(c)} + A_{\mathrm{P}}^{(c)} - A_{\mathrm{P}}^{(t)}\right] \tag{4}$$

and

$$ae^{i\theta} \equiv R_b \left[\frac{A_{\rm P}^{(u)} - A_{\rm P}^{(t)}}{A_{\rm T}^{(c)} + A_{\rm P}^{(c)} - A_{\rm P}^{(t)}} \right] \tag{5}$$

are CP-conserving, "hadronic" parameters, where

$$A \equiv \frac{|V_{cb}|}{\lambda^2} = 0.809 \pm 0.026 \quad \text{and} \quad R_b \equiv \left(1 - \frac{\lambda^2}{2}\right) \frac{1}{\lambda} \left| \frac{V_{ub}}{V_{cb}} \right|, \tag{6}$$

with $\lambda \equiv |V_{us}| = 0.22521 \pm 0.00083$ are CKM parameters. It should be noted that R_b is one of the sides of the UT.

The $B_s^0 \to J/\psi K_{\rm S}$ channel is a decay into a CP eigenstate with eigenvalue -1, and offers a time-dependent CP asymmetry of the following structure:

$$a_{\rm CP}(t) \equiv \frac{\Gamma(B_s^0 \to J/\psi K_{\rm S}) - \Gamma(\bar{B}_s^0 \to J/\psi K_{\rm S})}{\Gamma(B_s^0 \to J/\psi K_{\rm S}) + \Gamma(\bar{B}_s^0 \to J/\psi K_{\rm S})} =$$

$$\frac{C(B_s^0 \to J/\psi K_{\rm S}) \cos(\Delta M_s t) - S(B_s^0 \to J/\psi K_{\rm S}) \sin(\Delta M_s t)}{\cosh(\Delta\Gamma_s t/2) - \mathcal{A}_{\Delta\Gamma}(B_s^0 \to J/\psi K_{\rm S}) \sinh(\Delta\Gamma_s t/2)}, \tag{7}$$

where $C(B_s^0 \to J/\psi K_{\rm S})$ and $S(B_s^0 \to J/\psi K_{\rm S})$ describe direct and mixing-induced CP violation, respectively, and $\mathcal{A}_{\Delta\Gamma}(B_s^0 \to J/\psi K_{\rm S})$ is accessible thanks to the width difference $\Delta\Gamma_s$ of the B_s-meson system. In terms of CP-violating phases and the hadronic parameters introduced above, these observables read as follows:

$$C(B_s^0 \to J/\psi K_{\rm S}) = \frac{2a \sin\theta \sin\gamma}{1 - 2a\cos\theta\cos\gamma + a^2} \tag{8}$$

$$S(B_s^0 \to J/\psi K_{\rm S}) = \frac{\sin\phi_s - 2a\cos\theta \sin(\phi_s + \gamma) + a^2 \sin(\phi_s + 2\gamma)}{1 - 2a\cos\theta\cos\gamma + a^2} \tag{9}$$

$$\mathcal{A}_{\Delta\Gamma}(B_s^0 \to J/\psi K_{\rm S}) = \frac{\cos\phi_s - 2a\cos\theta \cos(\phi_s + \gamma) + a^2 \cos(\phi_s + 2\gamma)}{1 - 2a\cos\theta\cos\gamma + a^2}, \tag{10}$$

where ϕ_s is the CP-violating B_s^0–$\bar{B}_s^0$ mixing phase, which is in the Standard Model (SM) given by $\phi_s^{\rm SM} = -2\lambda^2\eta = -(2.12 \pm 0.11)^\circ$ but may well be enhanced by NP effects [7].

It should be noted that these quantities are not independent from one another, satisfying the relation

$$\left[C(B_s^0 \to J/\psi K_{\rm S})\right]^2 + \left[S(B_s^0 \to J/\psi K_{\rm S})\right]^2 + \left[\mathcal{A}_{\Delta\Gamma}(B_s^0 \to J/\psi K_{\rm S})\right]^2 = 1. \tag{11}$$

However, we have still another independent observable at our disposal, which is the CP-averaged rate [2]

$$\langle \Gamma(B_s^0 \to J/\psi K_{\rm S}) \rangle \equiv \Phi^s_{J/\psi K_{\rm S}} \times |\mathcal{N}|^2 \times \left[1 - 2a\cos\theta\cos\gamma + a^2\right], \tag{12}$$

where $\Phi^s_{J/\psi K_{\rm S}}$ denotes a phase-space factor, and $\mathcal{N} \equiv -\lambda\mathcal{A}$.

In the case of the decay $B_d^0 \to J/\psi K_{\rm S}$, the counterpart of the penguin parameter $ae^{i\theta}$ enters with the doubly Cabibbo-suppressed parameter $\epsilon \equiv \lambda^2/(1-\lambda^2) = 0.053$ as follows:

$$A(B_d^0 \to J/\psi\, K_{\rm S}) = \left(1 - \frac{\lambda^2}{2}\right) \mathcal{A}' \left[1 + \epsilon a' e^{i\theta'} e^{i\gamma}\right], \tag{13}$$

where the hadronic parameters are defined in analogy to the $B_s^0 \to J/\psi K_{\rm S}$ case [2], and the primes remind us that we are dealing with a $\bar{b} \to \bar{s}$ mode. The corresponding observables can be obtained from the expressions given above by simply making the following replacements:

$$a \to -a', \quad \theta \to \theta', \quad \mathcal{A} \to \mathcal{A}', \quad \mathcal{N} \to \mathcal{N}' \equiv -\frac{1}{\sqrt{\epsilon}}\frac{\mathcal{A}'}{\mathcal{A}}\mathcal{N}. \tag{14}$$

Concerning the mixing-induced CP asymmetry $S(B_d^0 \to J/\psi K_{\rm S})$, we have to replace, in addition, the B_s^0–$\bar{B}_s^0$ mixing phase by its B_d-meson counterpart ϕ_d, which is in the SM given by $\phi_d^{\rm SM} = 2\beta$.

Thanks to the ϵ suppression of the penguin parameter in (13), the $B_d^0 \to J/\psi K_{\rm S}$ decay is referred to as the "golden" channel for the measurement of ϕ_d. In the B-factory era of the last decade, it was justified in terms of accuracy to simply neglect this term. On the other hand, we are entering a new territory for precision B physics in the LHC era which makes it mandatory to take the penguin effects into account. This was already anticipated in 1999 in Ref. [2], where it was proposed to use the $B_s^0 \to J/\psi K_{\rm S}$ channel to accomplish this task and to extract the angle γ of the UT, exploiting the feature that the decays $B_s^0 \to J/\psi K_{\rm S}$ and $B_d^0 \to J/\psi K_{\rm S}$ are related to each other through an interchange of all down and strange quarks, i.e. by the U-spin flavour symmetry of strong interactions. The relevant observables are the CP asymmetries $C(B_s^0 \to J/\psi K_{\rm S})$ and $S(B_s^0 \to J/\psi K_{\rm S})$ as well as the following ratio of the CP-averaged rates:

$$H \equiv \frac{1}{\epsilon}\frac{\langle\Gamma(B_s^0 \to J/\psi K_{\rm S})\rangle}{\langle\Gamma(B_d^0 \to J/\psi K_{\rm S})\rangle}\left|\frac{\mathcal{A}'}{\mathcal{A}}\right|^2\frac{\Phi^d_{J/\psi K_{\rm S}}}{\Phi^s_{J/\psi K_{\rm S}}} = \frac{1-2a\cos\theta\cos\gamma+a^2}{1+2\epsilon a'\cos\theta'\cos\gamma+\epsilon^2 a'^2}. \tag{15}$$

The U-spin flavour symmetry implies

$$a = a', \quad \theta = \theta', \tag{16}$$

as well as

$$\mathcal{A}' = \mathcal{A}. \tag{17}$$

The three observables depend then on γ as well as a and θ. Consequently, we can convert the measured values of $C(B_s^0 \to J/\psi K_{\rm S})$, $S(B_s^0 \to J/\psi K_{\rm S})$ and H into γ, a and θ. In Sections 5 and 6, we discuss in detail how the implementation of this strategy works in practise at the LHCb experiment.

3 Picture from Current Data

The CDF collaboration has recently reported the observation of the $B_s^0 \to J/\psi K_{\rm S}$ channel, with the following result [1]:

$$\frac{{\rm BR}(B_s^0 \to J/\psi K_{\rm S})}{{\rm BR}(B_d^0 \to J/\psi K_{\rm S})} = 0.0405 \pm 0.0070({\rm stat.}) \pm 0.0041({\rm syst.}) \pm 0.0050({\rm frag.}), \tag{18}$$

where the last error is associated with the ratio f_s/f_d of the B_s and B_d fragmentation functions. Using the value $\mathrm{BR}(B^0_d \to J/\psi K^0) = (8.71 \pm 0.32) \times 10^{-3}$ of the Particle Data Group (PDG) [10] yields

$$\mathrm{BR}(B^0_s \to J/\psi \bar{K}^0) = (3.53 \pm 0.61(\mathrm{stat.}) \pm 0.35(\mathrm{syst.}) \pm 0.43(\mathrm{frag.}) \pm 0.13(\mathrm{PDG})) \times 10^{-5}. \tag{19}$$

It is interesting to compare this result with the branching ratio of the $B^0_d \to J/\psi\pi^0$ channel. If we neglect penguin annihilation and exchange topologies, which contribute to $B^0_d \to J/\psi\pi^0$ but have no counterpart in $B^0_s \to J/\psi\bar{K}^0$ and are expected to play a minor role (which can be probed through $B^0_s \to J/\psi\pi^0$), the $SU(3)$ flavour symmetry applied to the spectator d and s quarks implies

$$\Xi_{SU(3)} \equiv \frac{\mathrm{BR}(B^0_s \to J/\psi\bar{K}^0)}{2\mathrm{BR}(B^0_d \to J/\psi\pi^0)} \frac{\tau_{B_d}}{\tau_{B_s}} \frac{\Phi^d_{J/\psi\pi^0}}{\Phi^s_{J/\psi K_\mathrm{S}}} \xrightarrow{SU(3)} 1, \tag{20}$$

where the factor of 2 is associated with the wave function of the π^0, whereas the τ_{B_d} and τ_{B_s} denote the B_d and B_s lifetimes, respectively. Using the PDG values $\tau_{B_d}/\tau_{B_s} = 1.036 \pm 0.018$ and $\mathrm{BR}(B^0_d \to J/\psi\pi^0) = (1.76 \pm 0.16) \times 10^{-5}$, we obtain

$$\Xi_{SU(3)} = 1.01 \pm 0.25, \tag{21}$$

which agrees very well with the $SU(3)$ expectation in (20), although the error is still sizable.

In the following analysis, we will use the hadronic parameters extracted from the CP violation data for the $B^0_d \to J/\psi\pi^0$ channel as a guideline, using the numerical values obtained in Ref. [3]. Making the same assumptions as in (20), we have then

$$a \in [0.15, 0.67], \quad \theta \in [174, 213]^\circ, \tag{22}$$

which correspond to $\gamma = (65 \pm 10)^\circ$. The numerical value in (21) gives us confidence in this procedure. The observables of the $B^0_s \to J/\psi K_\mathrm{S}$ channel then read as

$$H = 1.53 \pm 0.31, \tag{23}$$

and

$$C \equiv C(B^0_s \to J/\psi K_\mathrm{S}) = C(B^0_d \to J/\psi\pi^0) = -0.10 \pm 0.13. \tag{24}$$

Concerning the mixing-induced CP violation $S \equiv S(B^0_s \to J/\psi K_\mathrm{S})$, we can calculate this observable with the help of Eq. (9) as a function of the B^0_s–$\bar{B}^0_s$ mixing phase ϕ_s for given values of H, C and γ. The corresponding formulae are given as follows:

$$2a\cos\theta = \frac{1 - H + (1 - \epsilon^2 H)a^2}{(1 + \epsilon H)\cos\gamma} \tag{25}$$

$$1 - 2a\cos\theta\cos\gamma + a^2 = \left[\frac{(1+\epsilon)(1+\epsilon a^2)}{1+\epsilon H}\right] H, \tag{26}$$

where

$$a^2 = P \pm \sqrt{P^2 - Q} \tag{27}$$

with

$$P=\frac{2\left[(1+\epsilon H)\sin\gamma\cos\gamma\right]^2-(1-H)(1-\epsilon^2H)\sin^2\gamma-\epsilon\left[(1+\epsilon)HC\cos\gamma\right]^2}{\left[(1-\epsilon^2H)\sin\gamma\right]^2+\left[\epsilon(1+\epsilon)HC\cos\gamma\right]^2} \tag{28}$$

$$Q=\frac{\left[(1+\epsilon)HC\cos\gamma\right]^2+\left[(1-H)\sin\gamma\right]^2}{\left[(1-\epsilon^2H)\sin\gamma\right]^2+\left[\epsilon(1+\epsilon)HC\cos\gamma\right]^2}. \tag{29}$$

In these expressions, we have used the U-spin relation (16) but did not make any other approximation. If we assume the SM, we arrive at the following prediction:

$$S(B_s^0\to J/\psi K_{\rm S})|_{\rm SM}=0.54\,{}^{+\,0.14}_{-\,0.25}|_H\,{}^{+\,0.00}_{-\,0.01}|_C\,{}^{+\,0.16}_{-\,0.13}|_\gamma=0.54\,{}^{+\,0.21}_{-\,0.28}. \tag{30}$$

As we have noted above, NP may well enter the $B_s^0\to J/\psi K_{\rm S}$ channel through contributions to B_s^0–$\bar{B}_s^0$ mixing, yielding a value of ϕ_s different from $\phi_s^{\rm SM}$. The most recent compilation of the CDF and DØ collaborations using measurements of CP violation in $B_s^0\to J/\psi\phi$ [11] can be found in Refs. [12] and [13], respectively. Unfortunately, the situation is not conclusive, although the CDF and DØ data are consistent with each other: the CDF collaboration finds the ranges $\phi_s\in[-59.6^\circ,-2.29^\circ]\sim-30^\circ\vee[-177.6^\circ,-123.8^\circ]\sim-150^\circ$ (68% C.L.), while the DØ collaboration gives a best fit value around $\phi_s\sim-45^\circ$, taking also information from the dimuon charge asymmetry and the measured $B_s^0\to D_s^{(*)+}D_s^{(*)-}$ branching ratio into account.

Interestingly, we can also probe this kind of NP through the mixing-induced CP violation in $B_s^0\to J/\psi K_{\rm S}$. In Fig. 2, we show the correlation between $S(B_s^0\to J/\psi K_{\rm S})$ and $\sin\phi_s$, which can be determined from the time-dependent angular analysis of the $B_s^0\to J/\psi\phi$ channel [11]. In this figure, we assume that NP is only present in B_s^0–$\bar{B}_s^0$ mixing and have used the formulae give above to calculate curves for different values of H and γ. We observe the following features:

- Thanks to the penguin contributions, we can distinguish between the twofold ambiguity for ϕ_s originating from the measured value of $\sin\phi_s$, in particular also between $\phi_s\sim0^\circ$ (as in the SM) and $\phi_s\sim180^\circ$.

- For $\phi_s\sim-30^\circ$ and $\phi_s\sim-45^\circ$, the mixing-induced CP asymmetry is ~0 while it approaches -1 for $\phi_s\sim-150^\circ$. Consequently, we could then also clearly identify the scenarios corresponding to the CDF and DØ data discussed above.

In view of these observations, already first experimental information on the mixing-induced CP violation in $B_s^0\to J/\psi K_{\rm S}$ would be very interesting to complement the analyses of CP violation in $B_s^0\to J/\psi\phi$. Needless to note that this feature relies on sizable penguin contributions to the $B_s^0\to J/\psi K_{\rm S}$ channel.

4 Framework for the LHCb Feasibility Study

With an anticipated integrated luminosity of 1 fb^{-1} by the end of 2011, the LHCb experiment expects to be able to measure $\sin\phi_s$ with a precision between 0.04 and



Figure 2: Correlation between $\sin\phi_s$ and $S(B_s^0 \to J/\psi K_{\rm S})$ for different values of H (left panel, with $\gamma = 65°$) and different values of γ (right panel, with $H = 1.53$). The impact of a variation of $C = -0.10 \pm 0.13$ is small.

0.07 [14]. The $B_s^0 \to J/\psi K_{\rm S}$ decay should then already be clearly visible with the following uncertainties:

$$\frac{{\rm BR}(B_s^0 \to J/\psi K_{\rm S})}{{\rm BR}(B_d^0 \to J/\psi K_{\rm S})} = 0.0403 \pm 0.0065({\rm stat.}) \pm 0.0029({\rm frag.}), \qquad (31)$$

based on an estimated event yield of 60 000 $B_d^0 \to J/\psi K_{\rm S}$ and 700 $B_s^0 \to J/\psi K_{\rm S}$ decays, and where the error on the fragmentation fractions is based on the method proposed in Ref. [15]. Concerning the CP-violating observables, we expect a sensitivity of $\Delta C = \Delta S = 0.34$, which might allow us to get first information on these quantities.

In order to explore the feasibility of the $B_s^0 \to J/\psi K_{\rm S}$ measurement at LHCb with higher integrated luminosities, we perform a toy Monte Carlo (MC) analysis. For this study we assume the following situation: Already with an integrated luminosity of $1\,{\rm fb}^{-1}$ we can pin down the B_s^0–$\bar{B}_s^0$ mixing phase. Hence, for the two milestone scenarios it will be well known. In the following discussion we will consider the SM case $\phi_s = -2.12°$. Secondly, the used signal and background yields are extrapolated from Ref. [5]. The number of $B_d^0 \to J/\psi K_{\rm S}$ events listed there is scaled, using the measured $B_s^0 \to J/\psi K_{\rm S}$ branching fraction, to give an expected yield of 4000 and 70 000 $B_s^0 \to J/\psi K_{\rm S}$ events for the integrated luminosities of 6 ${\rm fb}^{-1}$ and 100 ${\rm fb}^{-1}$, respectively. These event yields are based on an estimated inclusive J/ψ cross section of $\sigma_{{\rm Incl.}J/\psi} = 262\,\mu{\rm b}$ at $\sqrt{s} = 14$ TeV, the LEP fragmentation functions published by HFAG [16], and a $b\bar{b}$ cross section at $\sqrt{s} = 14$ TeV of $\sigma_{b\bar{b}} = 568\,\mu{\rm b}$, which was extrapolated from the recently measured $b\bar{b}$ cross section of $\sigma_{b\bar{b}} = (284 \pm 20\,({\rm stat.}) \pm 49\,({\rm syst.}))\,\mu{\rm b}$ at $\sqrt{s} = 7$ TeV [17].

For the MC analysis the reconstructed mass and lifetime distribution of both signal and background are modelled. The former is used to obtain the errors on the CP observables C and S, while the latter is used for the determination of the statistical error on H. The error on H coming from the ratio f_s/f_d of fragmentation functions,

Parameter	Value	Parameter	Value
$\sigma_{b\bar{b}}$	568 μb	B_s lifetime resolution	0.041 ps
$\sigma_{\text{Incl.}J/\psi}$	262 μb	mistag fraction	0.36
f_d/f_s	3.55 ± 0.26	B mass resolution	15.2 MeV/c^2

Table 1: Overview of the parameters used by the toy MC.

affecting H via (18), is again determined using the strategy proposed in Ref. [15]. The most import parameters used by the toy MC to describe both distributions are listed in Table 1.

The modelled B_s lifetime distribution, given in the left panels of Fig. 3, has three major components: the $B_s^0 \to J/\psi K_{\text{S}}$ signal (plotted for B_s and $\bar{B}_s$ combined), an inclusive J/ψ background and a combinatoric $b\bar{b}$ background. The B_s and $\bar{B}_s$ signals are described individually using the time evolution equations (11) and (12) in Ref. [2], respectively, and include a dilution factor due to mistagging of the original B meson. The inclusive J/ψ sample contains both prompt and non-prompt (e.g. from B decays) J/ψ candidates, which results in an asymmetric distribution. The combinatoric $b\bar{b}$ background describes all other background from $b\bar{b}$ events; other sources of background are considered negligible. All lifetime distributions are convoluted with a Gaussian resolution model, which causes some events to have a negative lifetime. The resolution for the $B_s^0 \to J/\psi K_{\text{S}}$ signal is $\sigma = 0.041$ ps [5].

The modelled mass distribution, also given in Fig. 3, describes the two B mass peaks (B_d on the left, and B_s on the right and in the insert) and an exponential background for both the inclusive J/ψ and combinatoric $b\bar{b}$ sample.

The signal yield is small compared to the number of background events, as can be seen from the inserts in Fig. 3, making it hard to distinguish the B_s mass peak from the background. This is the main reason for choosing the high luminosity scenarios of 6 fb^{-1} and 100 fb^{-1}. Nonetheless, the lifetime distribution shows that the background is predominantly prompt. The long lifetime region ($t > 2$ ps) is therefore sufficiently sensitive to $B_s^0 \to J/\psi K_{\text{S}}$ in order to extract C and S. This is confirmed by the plot of the time-dependent CP asymmetry $a_{\text{CP}}(t)$ defined in equation (7) and given in Fig. 4 for our two milestone scenarios.

The toy MC study is mainly used to estimate the errors on the relevant observables. The actual values of these observables are determined by their dependency on a, θ and γ, given in equations (8), (9) and (15). The input values for a, θ and γ were taken from the $B_d^0 \to J/\psi\pi^0$ analysis whose result was already given in (22). We take the central values of these intervals,

$$a = 0.41 \quad , \quad \theta = 194^\circ \quad , \quad \gamma = 65^\circ, \tag{32}$$

which result in the following picture for the statistical uncertainties at LHCb:

$$6 \text{ fb}^{-1} : \Delta H = 0.069, \;\; \Delta S = 0.14, \quad \Delta C = 0.14, \tag{33}$$

$$100 \text{ fb}^{-1} : \Delta H = 0.052, \;\; \Delta S = 0.035, \;\; \Delta C = 0.035. \tag{34}$$

In principle these statistical errors could depend on the input value for a, θ and γ, but we will assume them constant for the remainder of this paper.



Figure 3: The lifetime distribution for an originally tagged B_s meson (left) and the total mass distribution (right) for the decay $B_s^0 \to J/\psi K_S$ and its background, as used in the MC analysis. This shows good sensitivity to the signal in the long lifetime region. The number of events correspond to a data sample of 6 fb^{-1} (top) and 100 fb^{-1} (bottom) respectively.



Figure 4: Toy MC simulation of the time-dependent CP asymmetry $a_{\mathrm{CP}}(t)$ defined in equation (7) for the $6\,\mathrm{fb}^{-1}$ (top) and $100\,\mathrm{fb}^{-1}$ (bottom) scenarios.

For a more detailed discussion of the experimental feasibility study, the reader is referred to Ref. [18].

5 Extraction of γ at LHCb

First information about γ could in principle be obtained from H as the general structure of this observable in (15) implies the following inequality [19]:

$$H \geq \left[1 - 2\epsilon \cos^2\gamma + \mathcal{O}(\epsilon^2)\right] \sin^2\gamma \tag{35}$$

However, as the experimental data indicate a value of $H > 1$, the corresponding bound on γ is not effective.

As discussed in Ref. [2], using the B_s^0–$\bar{B}_s^0$ mixing phase as an input, we can convert the $B_s^0 \to J/\psi K_S$ CP asymmetries C and S into a contour in the γ–a plane, which is theoretically clean. Using the U-spin symmetry, we can determine another contour from H and $S(B_s^0 \to J/\psi K_S)$. Thanks to the U-spin relation in (16), the intersection of these curves allows us then to extract γ and the penguin parameter a. Knowing these parameters, the CP-conserving strong phase θ can be extracted as well. In Fig. 5, we illustrate the corresponding situation in the γ–a plane for our LHCb study.



Figure 5: Determination of γ and a from intersecting contours. Situation resulting from our LHCb feasibility study for 6 fb^{-1} (left) and 100 fb^{-1} (right).

The angle γ can also be extracted, together with a and θ, from a simultaneous χ^2 fit to C, S and H using the expressions in (8), (9) and (15). The corresponding numerical results read as follows:

$$6\,\text{fb}^{-1}: \quad \gamma = (65 \pm 10)^\circ, \quad a = 0.410 \pm 0.079, \quad \theta = (194 \pm 17)^\circ, \tag{36}$$

$$100\,\text{fb}^{-1}: \quad \gamma = (65.0 \pm 3.2)^\circ, \quad a = 0.410 \pm 0.023, \quad \theta = (194.0 \pm 4.1)^\circ. \tag{37}$$

Here we give only the statistical errors, which show that LHCb should have sensitivity to extract γ from the U-spin-related $B_{s,d} \to J/\psi K_S$ decays. Using other avenues to extract

γ, in particular theoretically clean strategies employing $B_s^0 \to D_s^{\mp} K^{\pm}$ and similar modes, LHCb expects the following accuracies for γ:

$$6\ \mathrm{fb}^{-1}: \quad \gamma = (65.0 \pm 3.1)^\circ \ \ [14], \quad 100\ \mathrm{fb}^{-1}: \quad \gamma = (65.0 \pm 1.0)^\circ \ \ [6], \tag{38}$$

which should be compared with the current situation emerging from $B \to D^{(*)} K^{(*)}$ analyses:

$$\gamma = (71^{+21}_{-25})^\circ \text{ (CKMfitter [20])}, \quad (78 \pm 12)^\circ \text{ (UTfit [21])}. \tag{39}$$

Consequently, LHCb should make significant progress with the determination of γ with respect to the unsatisfactory current situation.

Concerning the $B_{s,d} \to J/\psi K_{\mathrm{S}}$ strategy, we find a good sensitivity for this angle in our feasibility study. We have also to deal with theoretical uncertainties originating from U-spin-breaking effects. As far as the corrections to (16) are concerned, they actually play a very minor role for the determination γ as the a' terms in (15) are strongly suppressed by the tiny ϵ parameter in [2]; numerically, they give a correction for γ well below 1°. However, U-spin-breaking effects enter also through the ratio $|\mathcal{A}'/\mathcal{A}|$, which affects the determination of H as defined in (15). We find that an uncertainty of 10% for the relevant form-factor ratio $F_{B_d^0 K^0}(M_{J/\psi}^2; 1^-)/F_{B_s^0 \bar{K}^0}(M_{J/\psi}^2; 1^-)$, as considered in Ref. [3], corresponds to an error of about $\Delta\gamma|_{\mathrm{FF}} = 12^\circ$, which essentially matches the statistical error of 10° for our 6 fb^{-1} scenario in (36). For an LHCb upgrade with 100 fb^{-1}, a form-factor uncertainty of 2.5% would be required in order to match (37), which serves as the ultimate benchmark for future calculations of this quantity. We hope that already by the end of the second LHCb data taking period, i.e. 2014–15, we will have a better understanding of $SU(3)$ breaking through studies of various other U-spin- and $SU(3)$-related B-meson decays, as well as through improved theoretical studies, including in particular also non-factorisable U-spin-breaking corrections to $|\mathcal{A}'/\mathcal{A}|$. The current data from the Tevatron do not indicate sizable non-factorisable $SU(3)$-breaking effects [22], which is also supported by (21).

An important aspect of the determination of γ through the $B_{s,d} \to J/\psi K_{\mathrm{S}}$ strategy is that it may reveal a NP contribution to the $B_s^0 \to J/\psi K_{\mathrm{S}}$ decay amplitude. This would be indicated by a value of γ in conflict with the clean determinations through the pure "tree" decays summarised in (38). In the following discussion, we assume that we have negligible NP effects of this kind, i.e. assume the SM expressions for the decay amplitudes in (3) and (13).

6 Clean Determination of the $B_s^0 \to J/\psi K_{\mathrm{S}}$ Penguin Parameters at LHCb

The major application of the $B_s^0 \to J/\psi K_{\mathrm{S}}$ channel at LHCb will be the extraction of the hadronic penguin parameters (a, θ) and to take them into account in the determination of the B_d^0–$\bar{B}_d^0$ mixing phase ϕ_d from $(\sin 2\beta)_{J/\psi K_{\mathrm{S}}}$. Since our goal is to minimise here the U-spin-breaking corrections, we shall refrain from using the H observable and will use γ as given in (38) as an input. Following Ref. [3], we can then convert the CP asymmetries S and C into theoretically clean contours in the θ–a plane; their intersection allows us



Figure 6: Determination of a and θ from intersecting contours. Situation resulting from our LHCb feasibility study for 6 fb^{-1} (left) and 100 fb^{-1} (right).

then to extract a and θ. In Fig. 6, we illustrate the corresponding situation arising from our LHCb feasibility study. In order to guide the eye, we show also the contour corresponding to H, which would be interesting to obtain insights into U-spin-breaking effects. Performing again a χ^2 fit yields the following results:

$$6\ \text{fb}^{-1} : a = 0.41 \pm 0.14\,(\text{stat.}) \pm 0.02\,(\gamma)\,, \quad \theta = [194 \pm 16\,(\text{stat.}) \pm 0.8\,(\gamma)]^\circ\,, \tag{40}$$

$$100\ \text{fb}^{-1} : a = 0.41 \pm 0.03\,(\text{stat.}) \pm 0.01\,(\gamma)\,, \quad \theta = [194.0 \pm 4.0\,(\text{stat.}) \pm 0.3\,(\gamma)]^\circ\,. \tag{41}$$

We observe that LHCb is expected to be able to determine the $B_s^0 \to J/\psi K_{\rm S}$ penguin parameters with impressive precision. It should be emphasised that these parameters rely only on the SM expression for the decay amplitude and are theoretically clean.

7 Controlling Penguin Effects in the Measurement of $(\sin 2\beta)_{J/\psi K_{\rm S}}$ at LHCb

The values of the penguin parameters a and θ are interesting from the point of view of strong interaction dynamics. However, they have also an important practical application for testing the Kobayashi–Maskawa mechanism of CP violation. Using the U-spin relation (16), we can relate the $B_s^0 \to J/\psi K_{\rm S}$ parameters to their $B_d^0 \to J/\psi K_{\rm S}$ counterparts, which allows us then to control the hadronic uncertainty of the measurement of the B_d^0–$\bar{B}_d^0$ mixing phase ϕ_d through the mixing-induced CP violation in $B_d^0 \to J/\psi K_{\rm S}$. The corresponding, generalised expression reads as follows [3]:

$$\frac{S(B_d^0 \to J/\psi K_{\rm S})}{\sqrt{1 - C(B_d^0 \to J/\psi K_{\rm S})^2}} = \sin(\phi_d + \Delta\phi_d), \tag{42}$$



Figure 7: Determining the penguin correction $\Delta\phi_d$ to $\sin(2\beta)$ from intersecting contours. Situation for 6 fb^{-1} (left) and 100 fb^{-1} (right).

where

$$\sin \Delta\phi_d = \frac{2\epsilon a \cos\theta \sin\gamma + \epsilon^2 a^2 \sin 2\gamma}{N\sqrt{1 - C(J/\psi K_\mathrm{S})^2}} \tag{43}$$

$$\cos \Delta\phi_d = \frac{1 + 2\epsilon a \cos\theta \cos\gamma + \epsilon^2 a^2 \cos 2\gamma}{N\sqrt{1 - C(J/\psi K_\mathrm{S})^2}} \tag{44}$$

with $N \equiv 1 + 2\epsilon a \cos\theta \cos\gamma + \epsilon^2 a^2$, so that

$$\tan \Delta\phi_d = \frac{2\epsilon a \cos\theta \sin\gamma + \epsilon^2 a^2 \sin 2\gamma}{1 + 2\epsilon a \cos\theta \cos\gamma + \epsilon^2 a^2 \cos 2\gamma}. \tag{45}$$

The current value of the mixing-induced CP violation is given by

$$S(B_d^0 \to J/\psi K^0) = 0.655 \pm 0.024, \tag{46}$$

while the direct CP asymmetry is currently measured as

$$C(B_d^0 \to J/\psi K^0) = -0.003 \pm 0.020. \tag{47}$$

These numbers are averages of the BaBar and Belle data over the $B_d^0 \to J/\psi K_\mathrm{S}$ and $B_d^0 \to J/\psi K_\mathrm{L}$ final states, as obtained by the Heavy Flavour Averaging Group [16]. The value in (47) implies that the deviation of the square root in (42) from one is at most at the level of 0.0002, i.e. completely negligible.

Using these formulae, we can convert the contour plots in Fig. 6 into curves in the θ–$\Delta\phi_d$ plane, as shown in Fig. 7. In this analysis, we have uncertainties from U-spin-breaking corrections to (16), which we parametrise as follows:

$$a = \xi a' \quad , \quad \theta = \theta' + \Delta\theta. \tag{48}$$



Figure 8: Correlation between the hadronic shift $\Delta\phi_d$ and its error for our LHCb feasibility study for different input values of the penguin parameters a and θ. Situation for 6 fb^{-1} (left) and 100 fb^{-1} (right); note the different scales of the y axes.

If we vary $\xi \in [0.85, 1.15]$ and $\Delta\theta \in [-20°, 20°]$, and perform a χ^2 fit to C and S, we obtain

$$6\ \text{fb}^{-1} : \Delta\phi_d = \left[-2.23 \pm 0.78\ (\text{stat.}) \pm 0.12\ (\gamma)\ {}^{+0.29}_{-0.40}(\xi)\ {}^{+0.33}_{-0.07}(\Delta\theta)\right]^\circ, \tag{49}$$

$$100\ \text{fb}^{-1} : \Delta\phi_d = \left[-2.23 \pm 0.19\ (\text{stat.}) \pm 0.04\ (\gamma)\ {}^{+0.29}_{-0.40}(\xi)\ {}^{+0.33}_{-0.07}(\Delta\theta)\right]^\circ. \tag{50}$$

In order to study the dependence on the input values of the penguin parameters for our analysis, we show the correlation between the hadronic shift $\Delta\phi_d$ and its statistical error at LHCb in Fig. 8. The corresponding curves show nicely that we can actually well determine the $\Delta\phi_d$ correction for a wide range of (a, θ) that should contain the "true" values of these parameters. It should also be noted that already a small penguin contribution with $a = 0.1$ gives a correction of $\Delta\phi_d \sim -0.5°$.

In order to fully appreciate the results of this study, we have to look at the precision for $(\sin 2\beta)_{J/\psi K_\text{S}}$ at LHCb: we expect 0.022 for $2\,\text{fb}^{-1}$ [5], and 0.003–0.010 (depending on to be determined systematic errors) for $100\,\text{fb}^{-1}$ [6]. We extrapolate these numbers into 0.014 for 6 fb^{-1}, which should be compared with (46). For a central value of $S(B^0_d \to J/\psi K^0) \sim 0.655$, these numbers correspond to errors of about 1° and (0.2–0.8)° for the phase itself. In order to match these very impressive precisions, we definitely have to control the doubly Cabibbo-suppressed penguin contributions in $B^0_d \to J/\psi K_\text{S}$. Looking at the numerical results in (49) and (50), we observe that we can actually achieve this goal. Already with $6\,\text{fb}^{-1}$ we have to control the penguin effects, for an upgrade of LHCb this is absolutely necessary to fully exploit the tremendous statistics. Measurements along these lines may eventually allow us to resolve CP-violating NP contributions to B^0_d–$\bar{B}^0_d$ mixing.

Let us finally note that we can – in addition to the direct CP violation in the $B^0_d \to J/\psi K_{\text{S,L}}$ channels [3] – add another observable to this analysis. It is the direct

CP asymmetry of the $B_u^+ \to J/\psi\pi^+$ decay, which we get from the $B_s^0 \to J/\psi K_{\mathrm{S}}$ topologies in Fig. 1 by replacing the strange spectator quark by an up quark. The resulting contours in Fig. 7 are similar to those related to C. Already with 1 fb^{-1}, as expected by the end of 2011, we find from a feasibility study similar to that described above that the $B_u^+ \to J/\psi\pi^+$ mode should be clearly visible at LHCb, with a statistical error of $C(B_u^\pm \to J/\psi\pi^\pm) = 0.005$. However, the production asymmetry between B and $\bar{B}$ states at the LHC (which, unlike the Tevatron, is a *pp* collider) is expected to be non-negligible and will have to be measured and corrected for. This is likely to induce irreducible systematic errors at a few per-cent level for the measurement of $C(B_u^\pm \to J/\psi\pi^\pm)$, which have to be studied in further detail.

8 Conclusions

As was pointed out about one decade ago, the decay $B_s^0 \to J/\psi K_{\mathrm{S}}$ is the U-spin partner of the "golden" mode $B_d^0 \to J/\psi K_{\mathrm{S}}$ and allows an extraction of the UT angle γ and of the corresponding penguin parameters. In the summer of 2010, the CDF collaboration has announced the first observation of the $B_s^0 \to J/\psi K_{\mathrm{S}}$ transition, with a branching ratio in agreement with an $SU(3)$ relation to the $B_d^0 \to J/\psi\pi^0$ decay.

We have calculated a correlation between the mixing-induced CP violation in the $B_s^0 \to J/\psi K_{\mathrm{S}}$ channel, S, and $\sin\phi_s$, which serves as a "target region" for the first CP violation measurement in this decay. We observe an interesting pattern for values of the B_s^0–$\bar{B}_s^0$ mixing phase ϕ_s in the region of the current measurements from the Tevatron: we can distinguish between the twofold solutions for ϕ_s arising from the measurement of $\sin\phi_s$, with $S \sim 0$ and $S \sim -0.8$ for $\phi_s \sim -30^\circ$ and $\phi_s \sim -150^\circ$, respectively, whereas we expect $S \sim +0.5$ in the SM. Thanks to improved measurements of input parameters, these predictions can be sharpened in the future.

By the end of 2011, i.e. the end of the first LHCb data taking period, we expect a clear signal for $B_s^0 \to J/\psi K_{\mathrm{S}}$, with very crude first information on the corresponding CP asymmetries. Extrapolating from published LHCb studies, we have performed a detailed feasibility study for the measurement of the $B_s^0 \to J/\psi K_{\mathrm{S}}$ observables, yielding about 4000 and 70000 signal events for 6 fb^{-1} and 100 fb^{-1}, respectively. These integrated luminosities refer to the end of the second LHCb data taking period of 2014–15 and a subsequent LHCb upgrade. Using input parameters as suggested by current data, we have shown that the $B_{s,d} \to J/\psi K_{\mathrm{S}}$ strategy offers indeed another determination of γ for LHCb. However, we conclude that the most important application of this method will be the determination of the corresponding hadronic penguin parameters and their control in the measurement of $(\sin 2\beta)_{J/\psi K_{\mathrm{S}}}$, which will allow us to match the corresponding experimental precision. Studies along these lines may eventually allow us to resolve NP in B_d^0–$\bar{B}_d^0$ mixing.

Acknowledgements
We would like to thank Marcel Merk and Wouter Hulsbergen for carefully reading the manuscript and useful comments and suggestions.

References

[1] CDF Collaboration, CDF Note 10240, August 2010, see also http://www-cdf.fnal.gov/physics/new/bottom/100708.blessed-BsJpsiK/

[2] R. Fleischer, Eur. Phys. J. C **10** (1999) 299 [arXiv:hep-ph/9903455].

[3] S. Faller, M. Jung, R. Fleischer and T. Mannel, Phys. Rev. D **79** (2009) 014030 [arXiv:0809.0842 [hep-ph]].

[4] M. Ciuchini, M. Pierini and L. Silvestrini, Phys. Rev. Lett. **95** (2005) 221804.

[5] LHCb Collaboration, Report CERN-LHCC-2003-030 (2003).

[6] LHCb Collaboration, Expression of Interest CERN-LHCB-2008-019 (2008).

[7] P. Ball and R. Fleischer, Eur. Phys. J. C **48** (2006) 413 [arXiv:hep-ph/0604249].

[8] L. Wolfenstein, Phys. Rev. Lett. **51** (1983) 1945.

[9] A. J. Buras, M. E. Lautenbacher and G. Ostermaier, Phys. Rev. D **50** (1994) 3433 [arXiv:hep-ph/9403384].

[10] K. Nakamura et al. (Particle Data Group), J. Phys. G **37** (2010) 075021.

[11] I. Dunietz, R. Fleischer and U. Nierste, Phys. Rev. D **63** (2001) 114015 [arXiv:hep-ph/0012219]; A. S. Dighe, I. Dunietz and R. Fleischer, Eur. Phys. J. C **6** (1999) 647 [arXiv:hep-ph/9804253].

[12] CDF Collaboration, CDF/ANAL/BOTTOM/PUBLIC/10206, July 2010, see also http://www-cdf.fnal.gov/physics/new/bottom

[13] DØ Collaboration, DØ Note 6093-CONF, July 2010, see also http://www-d0.fnal.gov/Run2Physics/WWW/results/b.htm

[14] LHCb Collaboration, [arXiv:0912.4179 [hep-ph]].

[15] R. Fleischer, N. Serra and N. Tuning, Phys. Rev. D **82** (2010) 034038 [arXiv:1004.3982 [hep-ph]].

[16] E. Barberio *et al.* (Heavy Flavour Averaging Group), [arXiv:0704.3575 [hep-ex]]; for the most recent updates, see http://www.slac.stanford.edu/xorg/hfag.

[17] R. Aaij *et al.* (LHCb Collaboration), arXiv:1009.2731 [hep-ex].

[18] K. De Bruyn, Master thesis, University of Amsterdam, *in progress.*

[19] R. Fleischer and S. Recksiegel, Phys. Rev. D **71** (2005) 051501 [arXiv:hep-ph/0409137]; R. Fleischer and T. Mannel, Phys. Rev. D **57** (1998) 2752 [arXiv:hep-ph/9704423].

[20] CKMfitter Collaboration, http://ckmfitter.in2p3.fr/

[21] UTfit Collaboration, http://www.utfit.org/UTfit/

[22] R. Fleischer, Eur. Phys. J. C **52** (2007) 267 [arXiv:0705.1121 [hep-ph]].